

SMS in protein-protein communication

How Short Motif Sequences are read
and understood

Ora Schueler-Furman



Faculty of Medicine, Hebrew University, Jerusalem, ISRAEL

Albert Szent-Györgyi Lecture, Eötvös Loránd University, Budapest 31/5/2016

Albert Szent-Györgyi

- So what the school has to do, in the first place, is to make us **learn how to learn**, to whet our appetites for knowledge, to teach us the delight of doing a job well and the excitement of creativity, to teach us to love what we do, and to help us to find what we love to do.

From: Teaching and expanding the knowledge (1964) Science 3649:1278

Communication by SMS

- Short Message Service
- Short Motif Sequences = peptides
- enhance communication

Communication by SMS

Communication may be :

- short
- regulated

Herbie (ERBB4) 1/3/2016 15:00
Hey Chap, R U free?
Give me a ride? 

Herbie (ERBB4) 1/3/2016 20:00
... I'm w8ing ... 

Chap (YAP1) 1/3/2016 21:00
Yep, I got a space




WOW (WWOX)
1/3/2016 21:00
Herbie, dinner! 

Herbie (ERBB4) 1/3/2016 21:05
Sorry Chap - 2 l8 

Communication by SMS


Communication may be :

- short
- regulated




SMS mediate many interactions

- transient
- regulation
- trans / cis
- PTM
- sequence-motifs (**SMS**)
- **structure :**
 - details of binding
 - drug design



SMS mediate many interactions




* London, Raveh, Movshovitz-Attias, Schueler-Furman. *Proteins* (2010);
<http://rosie.rosettacommons.org/peptiderive>, Sedan & Marcu, *NAR* (2016)

London, Barak and Schueler-Furman.
Current Opinion Chemical Biology (2013)

Peptide-protein interactions - Aims


1. define structure of peptide-protein interaction

Rosetta FlexPepDock & Cluspro PeptiDock




2. predict binding affinity & specificity


Rosetta FlexPepBind



3. detect new interactions




4. put things into context



Model structure: FlexPepdock


Nawsad Alam
Orly Marcu

Nir London
Barak Raveh
Lior Zimmerman



The challenge of peptide binding

Nature's challenge



$$\Delta G_{\text{BIND}} = \Delta H - T\Delta S$$

- ✧ **small** interface
- ✧ **flexible** peptide


➤ how do peptides do it?




London, Movshovitz-Attias, Schueler-Furman (2010) Structure

Nature's solutions

- ① increase ΔH




- ② decrease ΔS



prearranged receptor

The challenge of peptide binding

Our challenge



$$\Delta G_{\text{BIND}} = \Delta H - T\Delta S$$



- ✧ **small** interface
- ✧ **flexible** peptide




➤ how do we model these interactions?

Nature's solutions

- ① increase ΔH




- ② decrease ΔS



prearranged receptor

The challenge of peptide binding

Our challenge



$$\Delta G_{\text{BIND}} = \Delta H - T\Delta S$$


- ✧ **small** interface
- ✧ **flexible** peptide



➤ how do we model these interactions?


Our solution

- ① Extensive optimization of interface



- ② Restrict sampling to relevant space


➤ focus on peptide




Rosetta infrastructure for structural modeling

Sampling and optimization

“Monte-Carlo search with energy
Minimization” MCM (Liu *et al.*)



Energy function
(full-atom/centroid side-chains)




FlexPepDock protocol
for flexible peptide
docking and redesign

Loop modeling



Fragment libraries




Raveh, *et al.* Proteins (2010)

Raveh *et al.* PLoSOne (2011)

London *et al.* Current Opinion in Structural Biology (2013)

Rosetta FlexPepDock protocol



FlexPepDock



<http://flexpepdock.furmanlab.cs.huji.ac.il>

10 outer cycles


attractive term

repulsive term

rigid body moves
(Monte-Carlo with
Minimization)

backbone 'small' and
'shear' moves
(Monte-Carlo with Minimization)

200 or more models




General:
tested on large benchmark




- Short peptides 5-15 aa
- High resolution < 2 Å
- Non redundant < 70% seq id

Rosetta FlexPepDock

accurate prediction of motif residues




backbone rearrangements preserve critical interactions



➤ try our server :
<http://flexpepdock.furmanlab.cs.huji.ac.il/>





Impact of FlexPepDock

- first dedicated peptide-docking protocol
- motivated development of many new approaches
- CAPRI (docking competition) looks at peptides


→ to appear soon: THE book on peptide docking



edited with Nir London




Extension of Rosetta FlexPepDock



FlexPepDock



<http://flexpepdock.furmanlab.cs.huji.ac.il>



Ab initio
FlexPepDock



10 outer cycles

attractive term

repulsive term

rigid body moves
(Monte-Carlo with
Minimization)

backbone 'small' and
'shear' moves
(Monte-Carlo with Minimization)

200 or more models

Simulation temperature

rigid body
moves
(Monte-Carlo)

backbone moves : fragment
insertion + random moves
(Monte-Carlo)

10000-50000 models

➤Full *ab initio*




Model binding partners: FlexPepBind

Nawsad Alam


Nir London
Lior Zimmerman
Michal Sperber

Experiments:
Noah Wolfson
Fierke lab, U Michigan




FlexPepBind applications

Farnesyl transferase




London *et al.* PLoS Comput Biol (2011)

Geranyl-geranyl transferase




Sperber *et al.*, *in preparation*

Bcl2-BH3




London *et al.* Biochemistry (2012)

HDAC8




Decipher the deacetylome




Alam *et al.* Structure (2016)


Lysine (de)acetylation



Lysine acetyl transferase





Can we find more substrates?



Lysine deacetylase

“Oh no, that is not all I can do”
said the Kat in the Hat...

Lysine deacetylation



HDAC8

- ✓ active without co-factors
- ✓ related to important diseases


Known substrates:
P53, SMC3, ERR α , ARID1A,
Cortactin 9, Inv(16)

FlexPepBind Framework

1

INPUT

- peptide-protein complex structure



- training set

Binders

GYKacFGC
GIKacWGC
GRKacFGC


Non-Binders

GWKacMGC
GMKacGGC
GKKacKGC


2

CALIBRATION

- model set
- adjust protocol



- optimize for:



3


PREDICTION

- large-scale application

RESULTS

- identify new protein targets from discovered peptide substrates


...ERFEMFRELNEALELDAQ
AGKEPGGSRAHSSHLK**SKKG**
QTSRHKKLMFKTEGP...




HDAC8 substrate hunt: input

1

INPUT




binding site surrounded
by loops



K_{Ac} side
chain
planar &
stacked

K_{Ac} backbone fixed by 2
hydrogen bonds to D101



K_{Ac} acetyl coordinated by Zn ion

2v5w: HDAC8 - K-acetylated p53-
derived peptide with coumarine tail

HDAC8 substrate hunt: input (II)

1

INPUT

Binders

GYKacFGC
GIKacWGC
GRKacFGC



Non-Binders

GWKacMGC
GMKacGGC
GKKacKGC

training set
 $GX_1\text{Kac}X_2\text{GC}$

X_2



test set
 $GRKacX_2X_3\text{GC}$

X_3



SAMDI*



* Self-Assembled monolayers for Matrix assisted laser Desorption / Ionization time-of-flight mass spectrometry


Gurard-Levin et al. ACS Chem Biol. (2010).

HDAC8 substrates: critical features



2

CALIBRATION


- model training set
- adjust protocol
- predict test set



optimize for:



reinforce critical features
of interaction




favorable binding \uparrow = \uparrow favorable catalysis


HDAC8 substrate hunt: optimize

2 CALIBRATION


- model training set
- adjust protocol
- predict test set



optimize for
binary separation



training set **GX₁KacX₂GC**



thresholds:


stringent <-18.0 TP: 35% FP: 4%

lenient <-16.5 TP: 91% FP: 19%


HDAC8 substrate hunt: validate

2 CALIBRATION


- model training set
 - adjust protocol
 - predict test set



optimize for
binary separation



Test set GRKacX₂X₃GC




Summary of validation


2

CALIBRATION


- model training set
- adjust protocol
- predict test set



optimize for
binary separation



same, as histogram



Prediction of new HDAC8 substrates

3


PREDICTION

- apply protocol to set of candidate sequences

RESULTS

- identify new protein targets from discovered peptide substrates

...ERFEMFRELNEALELKDAQ
AGKEPGGSRAHSSH~~LK~~**SKKG**
QSTSRHKKLMFKTEGP...




PhosphoSitePlus®
with grant support from NIH LINCs Program, Italian School of Medicine at Monza Sismi, NIH
from Cell Signaling TECHNOLOGY®

Home | ABOUT PHOSPHOSITE | USING PHOSPHOSITE | CURATION PROCESS | CONTACT

PhosphoSitePlus® (PSP) is an online systems biology resource providing comprehensive information and tools for the study of protein post-translational modifications (PTMs) including phosphorylation, ubiquitination, acetylation and methylation. See [About PhosphoSite](#) above for more information.


Please cite the following reference for this resource: Hornbeck PV, et al (2015) *PhosphoSitePlus, 2014: mutations, PTMs and recalibrations*. Nucleic Acids Res. 43:D512-20. [reprint]

A PROTEIN MODIFICATION RESOURCE




Hornbeck et al. Proteomics. (2004)


PhosphoSite scanning: ~20'000 reported acetylated sites




In vitro HDAC8 activity assay



Prediction & experiment correlate




Summary of validation



- ✓ good correlation across experiments
- ✓ stable thresholds
- ✓ strongest measured HDAC8 substrates
 - peptide reactive ($>1000 K_{cat}/K_M$)
 - expected protein reactivity: $\times 1000$

Summary



Who are
these
substrates?

1

Strongest HDAC8 target: ZNF318

peptide	$\log(K_{cat}/K_M)$	min. I_{sc}	protein
FG K1275 FSW	3.68	-20.0	Zinc finger protein 318


K1275: unstructured & accessible

- G1274R in breast cancer
- S1277 phosphorylated

ZNF318 represses Androgen Receptor

- related partners:
 - sirt6, sirt7 & HDAC2

reactive ≥ 3 reasonable ≥ 2 mild ≥ 1 poor < 1




Targets: good old neighbors!

peptide	log (K _{cat} /K _M)	min. I_sc	protein	
STPV K ₂₉₂ FISR	2.20	-17.23	CSRP2BP (HAT)	
VS K ₃₅₀ GPF	2.09	-17.5	KAT6A (MYST3)	
DH K ₆₀₄ TLY	1.18	-16.5	KAT6A	
DS K ₁₅₈₃ NAK	1.02	-17.1	CREB Binding Protein	




KAT acetylated for activation (@604)



- HDAC8 -> KAT6A
- sirt1 -> hMOF

KAT350 : in unstructured regions

- fine tuning?




reactive ≥ 3 reasonable ≥ 2 mild ≥ 1 poor < 1

More chromatin remodelers ...

peptide	log(Kcat/ Km)	min. l_sc	protein
KLIS K18o8 FDKL	2.87	-17.85	ARID1A (chromatin remodeling)
SKIQ K3579 QLDQ	1.51	-16.25	MLL2 (methyltransferase)

- cross-talk between different modifications ?




reactive ≥ 3 reasonable ≥ 2 mild ≥ 1 poor < 1


More cross-talk

peptide	$\log(K_{cat}/K_M)$	min. I_sc	protein
SF K55 YAW	3.07	-20.8	EF1 α 1
LG K1017 FRR	2.54	-19.1	La-related protein 1
IS K504 YDR	1.92	-18.6	Msh6

- EF1 α 1 - transcription/translation
- Larp1 - binds top5' RNA
 - K1017 adjacent to RNA BD
- Msh6 - DNA mismatch repair
 - K504 in periphery of DNA


reactive ≥ 3 reasonable ≥ 2 mild ≥ 1 poor < 1






New targets - summary

✓ found strongest known (peptide) HDAC8 substrates



✓ role of HDAC8 modifications


✧ *peripheral* fine-tuning



✧ extensive potential for crossregulation

✓ structure-based identification of substrates

✧ binding is the bottleneck for catalysis



✧ protocol works for **flexible & internal** sites



Why a structure-based approach ?

peptide		log (Kcat/Km)	min. I_sc
KLIS	K1808 FDKL	2.87	-17.85
STPV	K292 FISR	2.20	-17.23
KRILH	K687 LLQN	1.70	-17.02
RVIGAK	K106 DQY	1.80	-16.4
SKIQ	K3579 QLDQ	1.51	-16.25
LGDG	K387 MKS	0.66	-14.79
KLSG	K167 EING	0.99	-14.29
KLGG	K1087 QRAA	1.04	-14.19
TEIG	K54 TLAEK	0.61	-13.96

structure and proteomic information complement each other


Proteomic identification of HDAC8 substrates

- ✓ *in vivo* data
- ✧ many known targets missed
- ✧ weak targets (GKac preferred) -> bias?

Why a structure-based approach ?


peptide		log (Kcat/ Km)	min. I_sc
DI K42	YPL	1.20	-18.6
VS K350	GPF	2.09	-17.5
DS K1583	NAK	1.02	-17.1
SG K256	YDL	1.12	-17.1
VS K93	GTL	1.23	-16.6
SF K769	SDQ	1.39	-16.6
DH K604	TLY	1.18	-16.5
LS K747	FLR	2.80	-21.6
SF K55	YAW	3.07	-20.8
FG K1275	FSW	3.68	-20.0
TW K78	ANF	1.89	-19.6
LG K1017	FRR	2.54	-19.1
IS K504	YDR	1.92	-18.6
KI K31	RLR	1.58	-18.5
SL K269	EFY	1.89	-18.0
RL K713	YSQ	1.90	-16.8
SG K517	YYY	1.32	-16.8
SG K8	YFA	1.30	-16.2
TF K292	GVD	1.15	-15.4
KG K878	DAE	1.56	-14.5
GG K149	AFG	1.32	-14.5
RK K1970	GEP	1.00	-11.9

KLIS	K1808	FDKL	2.87	-17.85
STPV	K292	FISR	2.20	-17.23
KRILH	K687	LLQN	1.70	-17.02
RVIGAK	K106	DQY	1.80	-16.4
SKIQ	K3579	QLDQ	1.51	-16.25
KLGG	K1087	QRAA	1.04	-14.19





sequence based motif Kac[FY]

- ✓ identifies 6/7 strong substrates
- ✧ many false positives
- ✧ weak substrates missed




What next ?

- ✓ found strongest known (peptide) HDAC8 substrates



- biology of new targets & functional impact
- differences and similarities among HDAC substrate specificities
 - ❖ targeted drugs ?
- extend to new systems (lipids, sugars, nucleotides...)



Back to the context

Jamal Fahoum


Emma-Joy Dodson

Shahar Rotem

Jordan Chill
Keren Adamsky
NMR Bar Ilan


Rami Aqeilan
Saleh Khawaled

Reuven Wiener



Back to the challenge

Nature's challenge



$$\Delta G_{\text{BIND}} = \Delta H - T\Delta S$$

- ❖ **small** interface
- ❖ **flexible** peptide


➤ how do peptides do it?




London, Movshovitz-Attias, Schueler-Furman (2010) Structure

Nature's solutions

- ① increase ΔH




- ② decrease ΔS



prearranged
receptor
& peptide

Back to the challenge

Nature's challenge




$$\Delta G_{\text{BIND}} = \Delta H - T\Delta S$$

- ❖ small interface
- ❖ flexible peptide

Nature's solutions

③ multiplicity




➤ opportunities for regulation by context!

Communication by SMS

When Wone is not enough

- many WW domains occur *in tandem*






Communication by SMS

experimental system :

- WWOX, YAP, ErbB4

techniques :

- ITC, CD, NMR, X-ray
- *in vitro* evolution




When None is not enough

50% of WW domains occur *in tandem*

- 👉 context crucial for complex regulation
- 👉 variable integration strategies

Dodson,..., Schueler-Furman (2015). Versatile communication strategies among tandem WW domain repeats. *Experimental Biology and Medicine*, 240:351



THANKS!

Fierke Lab U Michigan

Noah Wolfson
Caleb Joseph
Carol Fierke

Current lab members:

Nawsad Alam, Orly Marcu, Emma-Joy Dodson, Jamal Fahoum, Adily Mashash, Yuval Sedan, Oriel Goldstein, Aviou Zini, Alon Lalazry, Debbie Shalev, Shahar Rotem

Previous lab members

Nir London (Weizmann Institute), Barak Raveh (UCSF), Dan Reshef Lavanya Pushpam, Michal Sperber, Vered Fishbain, Lior Zimmerman, Assaf Lavi, Dana Movshovitz-Attias (CMU)

Dima Kozakov

Boston U & SUNY Stony Brook

Rosetta Community



Furman
Lab @ HU



NMR Jordan Chill, Keren Adamsky, Bar Ilan University

ITC & CD Assaf Friedler, Daniel Harries

Crystallography Reuven Wiener

In vivo assays Rami Aqeilan, Saleh Khawaled

